

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, 15/82, A01H 5/00		A1	(11) International Publication Number: WO 99/38979 (43) International Publication Date: 5 August 1999 (05.08.99)
(21) International Application Number: PCT/US99/01815 (22) International Filing Date: 28 January 1999 (28.01.99) (30) Priority Data: 60/072,944 29 January 1998 (29.01.98) US (71) Applicant: DOW AGROSCIENCES LLC [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268 (US). (72) Inventors: PETOLINO, Joseph, F.; 270 Woodstock Court, Zionsville, IN 46077 (US). PAREDDY, Dayakar, R.; 665 Woodbine Drive East, Carmel, IN 46033 (US). HOPKINS, Nicole, L.; 2518 Chaseway Court, Indianapolis, IN 46268 (US). ARMSTRONG, Katherine; 11202 Bridlewood Trail, Zionsville, IN 46077 (US). (74) Agent: STUART, Donald, R.; Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: WHISKER-MEDIATED TRANSFORMATION OF PLANT CELL AGGREGATES AND PLANT TISSUES AND REGEN- ERATION OF PLANTS THEREOF			
(57) Abstract Plant cell aggregates and plant tissues can be transformed by elongated, needle-like structures called "whiskers". The process comprises the agitation of plant cell aggregates and plant tissues of the plant to be transformed in the presence of DNA and whiskers, whereby DNA uptake and integration thereof is facilitated. The process may be applicable to other plant cell aggregates and plant tissues which have not proven easily transformable by other techniques.			

Best Available Copy

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WHISKER-MEDIATED TRANSFORMATION OF PLANT CELL AGGREGATES
AND PLANT TISSUES AND REGENERATION OF PLANTS THEREOF

This invention relates to a method of using
5 elongated, needle-like microfibers or "whiskers" to
transform plant cell aggregates and selected plant
tissues.

Until recently, genetically manipulated plants were
limited almost exclusively to those events created by
10 application of classical breeding methods. Creation of
new plant varieties by breeding was reserved primarily for
the most agronomically important crops, such as corn, due
to the cost and time needed to identify, cross, and stably
fix a gene in the genome, thus creating the desired trait.
15 In comparison, the advent of genetic engineering has
resulted in the introduction of many different
heterologous genes and subsequent traits into diverse
crops including corn, cotton, soybeans, wheat, rice,
sunflowers and canola in a more rapid manner. However,
20 the intergression of a new transgene into elite germplasm
is still quite a laborious task due to the tissue
culturing and back-crossing needed to produce a
commercially viable, elite, line.

Several techniques exist which allow for the
25 introduction, plant regeneration, stable integration, and
expression of foreign recombinant vectors containing
heterologous genes of interest in plant cells. One such
technique involves acceleration of microparticles coated
with genetic material directly into plant cells (U.S.
30 Patents 4,945,050 to Cornell; 5,141,131 to DowElanco; and
5,538,877 and 5,538,880, both to Dekalb). This technique
is commonly referred to as "microparticle bombardment" or
"biolistics". Plants may also be transformed using

Agrobacterium technology (U.S. Patent 5,177,010 to University of Toledo, 5,104,310 to Texas A&M, European Patent Application 0131624B1, European Patent Applications 120516, 159418B1 and 176,112 to Schilperoot, U.S. Patents 5,149,645, 5,469,976, 5,464,763 and 4,940,838 and 4,693,976 to Schilperoot, European Patent Applications 116718, 290799, 320500 all to Max Planck, European Patent Applications 604662,627752 and US Patent 5,591,616 to Japan Tobacco, European Patent Applications 0267159, and 0292435 and U.S. Patent 5,231,019 all to Ciba-Geigy, U.S. Patents 5,463,174 and 4,762,785 both to Calgene, and U.S. Patents 5,004,863 and 5,159,135 both to Agracetus).

Another transformation method involves the use of elongated needle-like microfibers or "whiskers" to transform cell suspension cultures (U.S. Patents 5,302,523 and 5,464,765 both to Zeneca). In addition, electroporation technology has been used to transform plant cells from which fertile plants have been obtained (WO 87/06614 to Boyce Thompson Institute; 5,472,869 and 5,384,253 both to Dekalb; 5,679,558, 5,641,664, WO9209696 and WO9321335 to Plant Genetic Systems).

Despite all of the technical achievements, genetic transformation and routine production of transgenic plants in a commercially viable, elite, germplasm is still a laborious task. For example, microparticle bombardment, while capable of being used either on individual cells, cell aggregates, or plant tissues, requires preparing DNA-attached gold particles and optimization of an expensive and not yet widely available, "gun" apparatus. Techniques involving *Agrobacterium* are extremely limited because not all plant species or varieties within a given species are susceptible to infection by the bacterium.

Electroporation techniques are not preferred due to the

extreme difficulties and cost typically encountered in routinely making protoplast from different plant species and tissues thereof and the concomitant low viability and low transformation rate associated therewith.

5 As disclosed herein, applicants have invented a method whereby plant cell aggregates and plant tissues from non-elite and elite germplasm can be directly and inexpensively transformed with a recombinant vector containing the gene of choice using whiskers. Applicants' invention is advantageous over currently used methods in that it is simple, quick and easy to use. Furthermore, applicants' invention is superior to that described in the art in that it eliminates the need to establish Type III callus cultures or establish and maintain cell suspension cultures, and can be used with either Type I or Type II callus, thus, is less germplasm limited. This means that applicants' invention, as described herein, can be used to transform elite genotypes directly thus eliminating the problems and time generally associated with gene intergression.

20 The present invention relates to the production of fertile, transgenic, *Zea mays* plants containing heterologous DNA preferably integrated into the chromosome of said plant and heritable by the progeny thereof.

25 One aspect of the present invention relates to *Zea mays* plants, plant parts, plant cells, plant cell aggregates, and seed derived from transgenic plants containing said heterologous DNA.

30 The present invention also relates to the production of fertile, transgenic, *Oryza sativa* L. plants containing heterologous DNA preferably integrated into the chromosome of said plant and heritable by the progeny thereof.

Another aspect of the present invention relates to *Oryza sativa* L. plants, plant parts, plant cells, plant cell aggregates, and seed derived from transgenic plants containing said heterologous DNA.

5 The present invention also relates to the production of fertile, transgenic, *Gossypium hirsutum* L. plants containing heterologous DNA preferably integrated into the chromosome of said plant and heritable by the progeny thereof.

10 Another aspect of the present invention relates to *Gossypium hirsutum* L. plants, plant parts, plant fibers, plant cells, plant cell aggregates, and seed derived from transgenic plants containing said heterologous DNA.

 The invention further relates to a process for
15 producing fertile transformed plants from Type I callus, Type II callus, hypocotyl-derived callus, or cotyledon-derived callus by whisker-mediated transformation.

 The invention yet further relates to a process for producing fertile transformed plants from meristematic
20 tissue by whisker-mediated transformation.

 Another aspect of the invention relates to fertile, mature maize plants regenerated from Type I or Type II callus and transgenic seed produced therefrom.

 Another aspect of the invention relates to fertile,
25 mature rice plants regenerated from Type I callus and transgenic seed produced therefrom.

 Yet, another aspect of the invention relates to fertile, mature cotton plants regenerated from hypocotyl-derived callus or cotyledon-derived callus and transgenic
30 seed and fiber produced therefrom.

 In a preferred embodiment, this invention produces the fertile transgenic plants described herein by means of whisker-mediated cell perforation and heterologous DNA

uptake, said whisker-mediated cell perforation being performed on plant cell aggregates and plant tissues, followed by a controlled regimen for selection and production of transformed plant lines.

5 Other aspects, embodiments, advantages, and features of the present invention will become apparent from the following specification.

The present invention relates to methods for production of fertile transgenic plants and seeds of, for
10 example, the species *Zea mays*, *Oryza sativa* L., *Gossypium hirsutum* L., and *Brassica napus* by transforming plant cell aggregates and plant tissues of said species with the DNA construct of interest via whisker-mediated transformation. After transformation, transgenic plants are regenerated
15 from said transformed plant cell aggregates and plant tissues and said regenerated plants express the chimeric DNA construct of interest. The transgenic plants produced herein by the methods described include: all species of corn including but not limited to field corn, popcorn,
20 sweet corn, flint corn, dent corn and the like; all species of cotton; and all species of rice.

The following phrases and terms are defined below:

By "antisense" is meant an RNA transcript that comprises sequences complementary to a target RNA and/or
25 mRNA or portions thereof and that blocks the expression of a target gene by interfering with the processing, transport, and/or translation of its primary transcript and/or mRNA. The complementarity may exist with any part of the target RNA, i.e., the 5' non-coding sequence,
30 3' non-coding sequence, introns, or the coding sequence. Antisense RNA is typically a complement (mirror image) of the sense RNA.

By "cDNA" is meant DNA that is complementary to and derived from a mRNA.

By "chimeric DNA construction" is meant a recombinant DNA containing genes or portions thereof from one or more species in either the sense or antisense orientation.

By "constitutive promoter" is meant promoter elements that direct continuous gene expression in all cell types and at all times (i.e., actin, ubiquitin, CaMV 35S, 35T, and the like).

By "cosuppression" is meant the introduction of a foreign gene having substantial homology to an endogenous gene, and in a plant cell causes the reduction in activity of the foreign gene and/or the endogenous gene product. Cosuppression can be sometimes achieved by introducing into said plant cell either the promoter sequence, the 5' and/or 3' ends, introns or the coding region of a gene.

By "developmental specific" promoter is meant promoter elements responsible for gene expression at specific plant developmental stages, such as in early or late embryogenesis and the like.

By "enhancer" is meant nucleotide sequence elements which can stimulate promoter activity such as those from maize streak virus (MSV), alfalfa mosaic virus (AMV), alcohol dehydrogenase intron 1 and the like.

By "expression" as used herein, is meant the transcription of enzymatic nucleic acid molecules, mRNA, and/or the antisense RNA inside a plant cell. Expression of genes also involves transcription of the gene and may or may not involve translation of the mRNA into precursor or mature proteins.

By "foreign" or "heterologous gene" is meant a gene having a DNA sequence that is not normally found in the

host cell, but is introduced by whisker-mediated transformation.

By "gene" is meant to include all genetic material involved in protein expression including chimeric DNA constructions, genes, plant genes and portions thereof.

By "genome" is meant genetic material contained in each cell of an organism and/or virus.

By "inducible promoter" is meant promoter elements which are responsible for expression of genes in response to a specific signal, such as: physical stimuli (heat shock genes); light (RUBP carboxylase); hormone (Em); metabolites, stress and the like.

By "modified plant" is meant a plant wherein the mRNA levels, protein levels or enzyme specific activity of a particular protein have been altered relative to that seen in an unmodified plant. Modification can be achieved by methods such as antisense, cosuppression, or over-expression.

By "plant cell aggregates", "plant cell lines", and "callus cell lines" is meant proliferating masses of tissue composed of a combination of undifferentiated and differentiated cells undergoing *de novo* morphogenesis and formed by placing a piece of plant material (explant) onto a growth-supporting medium under sterile conditions. The terms "plant cell aggregates", "plant cell lines", and "callus cell lines" are meant to include Type I and Type II callus cultures in monocotyledonous plants and hypocotyl- and cotyledon derived cultures in dicotyledonous plants. The terms defined herein are not intended to include either plant cell suspension cultures or Type III callus cultures.

By "plant tissues" is meant organized tissues including but not limited to meristems, embryos, pollen, cotyledons, germ cells, and the like.

By "promoter regulatory element" is meant nucleotide
5 sequence elements within a nucleic acid fragment or gene which controls the expression of that nucleic acid fragment or gene. Promoter sequences provide the recognition for RNA polymerase and other transcriptional factors required for efficient transcription. Promoter
10 regulatory elements from a variety of sources can be used efficiently in plant cells to express sense and antisense gene constructs. Promoter regulatory elements are also meant to include constitutive promoters, tissue-specific promoters, developmental-specific promoters, inducible
15 promoters and the like. Promoter regulatory elements may also include certain enhancer sequence elements that improve transcriptional or translational efficiency.

By "tissue-specific" promoter is meant promoter elements responsible for gene expression in specific cell
20 or tissue types, such as the leaves or seeds (i.e., zein, oleosin, napin, ACP, globulin and the like).

By "transgenic" is meant to include any Type I or Type II callus, hypocotyl- or cotyledon derived callus, tissue, plant parts or plants which contains heterologous
25 DNA or a chimeric gene construct that was introduced into said callus, tissue, plant parts or plants by whiskers and was subsequently transferred to later generations by sexual or asexual cell crosses or cell divisions.

By "whiskers" is meant elongated needle-like bodies
30 capable of being produced from numerous substances as described in "The Condensed Chemical Dictionary, Seventh Edition, Ed. Arthur & Elizabeth Rose, Reinhold Publishing Corp., New York (1966). The invention is not meant to be

limited to the material from which the whiskers are made but instead is meant to define a needle-like shaped structure wherein said whisker is smaller than the cell for which it is intended to be used in the transformation thereof. It is within the scope of this invention for whiskers to be shaped in a manner whereby DNA entry into a cell is facilitated. It is also intended that the scope of said invention include any material having a needle-like shape, said needle-like shaped material being able to perforate a plant cell with or without cell walls and thus facilitate DNA uptake and plant cell transformation. It is also intended that the scope of this invention not include microinjection techniques, such as wherein a DNA molecule is inserted into a cell by passing said DNA through an orifice intrinsic to a needle, said needle being first inserted into said cell. Preferably, whiskers are metal or ceramic needle-like bodies, with those most preferred being made of either silicon carbide or silicon nitride and being 30x0.5 μm to 10x0.3 μm in size.

By "whisker-mediated transformation" is meant the facilitation of DNA insertion into plant cell aggregates and/or plant tissues by whiskers and expression of said DNA in either a transient or stable manner.

In producing plant cell lines, tissues of interest are aseptically isolated and placed onto solid initiation medium whereby processes associated with cell differentiation and specialization occurring in organized plant cell tissues are disrupted, thus resulting in said tissues becoming dedifferentiated. Typically, initiation medium is solidified by adding agar or the like because callus cannot be readily initiated in liquid medium. Media are typically based on the N6 salts of Chu et al., (1978, Proc. Symp. Plant Tissue Culture, Peking Press, p

43-56) being supplemented with sucrose, vitamins, minerals, amino acids, and in some cases, synthetic hormones. However, callus tissues can also proliferate on media derived from the MS salts of Murashige and Skoog, (1962 *Physiol. Plant.* 15: 473-497). Cultures are generally maintained in a dark, sterile environment at about 28° C.

Typically, plant cell lines are preferably derived from tissues found in juvenile leaf basal regions, immature tassels, hypocotyl tissue, and cotyledonary nodes. For maize and rice, plant cell lines which produce meristematic tissue can be used, with those from zygotic embryo tissue being most preferred. Tissues most preferred for producing said plant cell lines are isolated from developing maize ears 10 to 14 days after pollination and non-germinated rice seed. Hypocotyl and cotyledon-derived tissues from seedlings are most preferred for production of cotton plant cell lines.

After placing said tissues on solid medium, new meristems arise after several days to a few weeks from either the scutellar region, in the case of corn and rice, or from hypocotyl or cotyledonary tissue in the case of cotton. These new meristems produce undifferentiated parenchymatous cell aggregates without the structural order characteristic of the tissue from which they are derived. Plant cell aggregates lack any recognizable overall structure and contain only a limited number of the many different kinds of specialized cell types found in intact, organized plant tissues. Said aggregates have been classified into non-embryogenic and embryogenic depending on their regenerative capacity, mode of reproduction, and tissue morphology (Franz, 1988, Ph.D. Thesis, University of Wageningen, The Netherlands).

In corn and rice, non-embryogenic calli are comprised of soft, granular, translucent tissue consisting of elongated, vacuolated cells incapable of plant regeneration. Alternatively, embryogenic,

5 monocotyledonous calli, being capable of somatic embryogenesis and plant regeneration, exist in three distinct morphotypes: Type I; Type II; and Type III.

Type I callus consists of compact, nodular, slow-growing embryogenic callus which proliferates as a mixture
10 of complex tissues exhibiting shoot and/or scutellar-like structures (Phillips et al, 1988, In, Corn and Corn Improvement, pp 345-387). Said callus is characterized by a high degree of cellular differentiation, well developed vascular structures and has been referred to as compact
15 embryogenic callus (US Patent 5,641,664 to Plant Genetic Systems). Essentially all monocotyledonous plants have tissue from which Type I callus can be produced. Plant regeneration in Type I callus normally occurs either through organogenesis by elongation of meristems (Green
20 and Phillips, 1975, Crop Sci., 15:417-421) and/or through somatic embryogenesis from a well defined root-shoot axis (Vasil et al., 1984, Amer. J. Bot. 71:158-161). The origin of regenerated shoots in Type I callus is not always obvious and appears to take place via sub-epidermal
25 meristem formation (Franz and Schel, 1991, Can. J. Bot. 69:26-33).

Type II callus, which is not as common as Type I, consists of soft, friable, embryogenic cells and can only be generated from certain monocotyledonous genotypes
30 (Phillips et al, 1988, In, Corn and Corn Improvement, pp 345-387). It grows rapidly, contains little or no vascular elements and can be described as friable, embryogenic callus (US Patent 5,641,664 to Plant Genetics

Systems). A distinguishing feature of Type II callus is that it contains numerous globular somatic embryos attached to suspensor-like structures on its surface through which plant regeneration appears to progress in clearly identifiable stages (Franz, 1988, Ph.D. Thesis, University of Wageningen, The Netherlands).

Type III callus has only been most recently described. Said callus is formed only very rarely, is easily dispersed in liquid and does not have any distinct somatic embryos on its surface. It has also been described as "friable, non-mucilaginous" callus (Shillito et al., 1989, Bio/Technology 7:581-587) and consists predominately of undifferentiated tissues capable of regeneration via somatic embryogenesis. Type III callus is considered the ideal tissue for transformation and regeneration because cells thereof easily disassociate and disperse, and thus, readily form suspension cultures. This type of callus is most rare, distinct from Type II callus, and can only be produced by visually selecting and preferentially enriching for it at each sub-culture passage of Type II cultures (WO94/28148; Zeneca).

In cotton, as with other dicotyledonous plants, callus types are typically not defined as Type I, II, or III. Moreover, hypocotyl- and cotyledon-derived callus cultures from cotton have been classified into non-embryogenic and embryogenic depending on morphology and regenerative capacity (Shoemaker et al., 1986 Plant Cell Rep. 3:178-181). Non-embryogenic callus is comprised of a loose, friable mass of cells that does not exhibit a strong cytoplasm staining reaction and cannot be used to readily regenerate plants. However, embryogenic callus appears as a tightly compact, dense cytoplasmic mass of cells capable of plant regeneration via somatic

embryogenesis. Somatic embryos produced therefrom first appear as globular structures which gradually elongate and then begin to exhibit cotyledonary development.

Of the callus types disclosed herein, Type I and Type II callus cultures derived from monocotyledonous plants are preferred in the production and regeneration of plants. Transgenic maize plants generated via whisker-mediated transformation are most preferably made from either Type I or Type II cultures; whereas, Type I callus cultures are most preferred in the production and regeneration of transgenic rice plants. In addition, cotyledonary node derived and hypocotyl-derived cultures are preferred in the production of transgenic cotton plants, with cultures produced from cotyledon tissue being most preferred.

Shoot tips of plants, including maize, rice, and cotton, contain apical meristems where organ primordia form from apical initial and subepidermal cells (Steeves and Sussex, 1989, Patterns in Plant Development, Cambridge University Press). Meristems can be isolated, placed onto shoot multiplication medium, and induced to produce multiple shoots from which plants can be regenerated (Zhong et al., 1992, Planta 187:483-489). Meristems, either freshly isolated or precultured to initiate shoot multiplication, can serve as recipient tissues for whisker-mediated transformation as taught in the present invention. Shoot tips containing meristematic regions are preferably removed from developing embryos (Lowe et al., 1995, Bio/Technology 13:677-682) or germinating seedlings (Gould et al., 1991, Plant Physiol. 95:426-434), transformed using whiskers with or without an osmotic pretreatment, and placed onto shoot proliferation medium

containing a selection agent prior to plant regeneration (Zhong et al., 1996, Plant Physiol. 110:1097-1107).

The heterologous DNA used for transformation herein may be circular, linear, double-stranded or single-stranded. Generally, said DNA is a recombinant vector plasmid and contains coding regions therein which serve to promote expression of the heterologous gene of interest as well as provide a selectable marker whereby those tissues containing said gene can be identified. Preferably, these recombinant vectors are capable of stable integration into the plant genome where selection of transformed plant lines is made possible by having said selectable marker expression driven either by constitutive, tissue-specific, or inducible promoters included therein.

One variable present in a heterologous DNA is the choice of the chimeric gene. Chimeric genes, either in the sense or antisense orientation, are expressed in plant cells under control of a constitutive, tissue-specific, developmental, or inducible promoter and the like. Preference for a particular chimeric gene is at the discretion of the artisan; however, chimeric genes can be, but are not limited to, from plants, animals, or bacteria and the like and can be used to express proteins either not found in a non-transformed cell or found in a transformed cell. Chimeric genes can be also used for, but are not limited to, up-regulation or down-regulation of an endogenous gene of interest.

Another variable is the choice of a selectable marker. Preference for a particular marker is at the discretion of the artisan, but any of the following selectable markers may be used along with any other gene not listed herein which could function as a selectable marker. Such selectable markers include but are not

limited to aminoglycoside phosphotransferase gene of transposon Tn5 (Aph II) which encodes resistance to the antibiotics kanamycin, neomycin and G418, as well as those genes which encode for resistance or tolerance to
5 glyphosate; hygromycin; methotrexate; phosphinothricin (bialophos); imidazolinones, sulfonylureas and triazolopyrimidine herbicides, such as chlorsulfuron; bromoxynil, dalapon and the like.

In addition to a selectable marker, it may be
10 desirous to use a reporter gene. In some instances a reporter gene may be used with or without a selectable marker. Reporter genes are genes which are typically not present in the recipient organism or tissue and typically encode for proteins resulting in some phenotypic change or
15 enzymatic property. Examples of such genes are provided in K. Weising et al. Ann. Rev. Genetics, 22, 421 (1988), which is incorporated herein by reference. Preferred reporter genes include the beta-glucuronidase (GUS) of the uidA locus of *E. coli*, the chloramphenicol acetyl
20 transferase gene from Tn9 of *E. coli*, the green fluorescent protein from the bioluminescent jellyfish *Aequorea victoria*, and the luciferase genes from firefly *Photinus pyralis*. An assay for detecting reporter gene expression may then be performed at a suitable time after
25 said gene has been introduced into recipient cells. A preferred such assay entails the use of the gene encoding beta-glucuronidase (GUS) of the uidA locus of *E. coli* as described by Jefferson et al., (1987 Biochem. Soc. Trans. 15, 17-19) to identify transformed cells.

30 Another variable is a promoter regulatory element. In addition to plant promoter regulatory elements, promoter regulatory elements from a variety of sources can be used efficiently in plant cells to express heterologous

genes. For example, promoter regulatory elements of bacterial origin, such as the octopine synthase promoter, the nopaline synthase promoter, the mannopine synthase promoter; promoters of viral origin, such as the
5 cauliflower mosaic virus (35S and 19S), 35T (which is a re-engineered 35S promoter, see PCT/US96/1682; WO 97/13402 published April 17, 1997) and the like may be used. Plant promoter regulatory elements include but are not limited to ribulose-1,6-bisphosphate (RUBP) carboxylase small
10 subunit (ssu), beta-conglycinin promoter, phaseolin promoter, ADH promoter, heat-shock promoters and tissue specific promoters.

Other elements such as matrix attachment regions, scaffold attachment regions, introns, enhancers,
15 polyadenylation sequences and the like may be present and thus may improve the transcription efficiency or DNA integration. Such elements may or may not be necessary for DNA function, although they can provide better expression or functioning of the DNA by affecting
20 transcription, stability of the mRNA and the like. Such elements may be included in the DNA as desired to obtain optimal performance of the transformed DNA in the plant. Typical elements include but are not limited to Adh-intron 1, the alfalfa mosaic virus coat protein leader sequence,
25 the maize streak virus coat protein leader sequence, as well as others available to a skilled artisan.

Constitutive promoter regulatory elements may also be used thereby directing continuous gene expression in all cells types and at all times (e.g., actin, ubiquitin,
30 CaMV 35S, and the like). Tissue specific promoter regulatory elements are responsible for gene expression in specific cell or tissue types, such as the leaves or seeds

(e.g., zein, oleosin, napin, ACP, globulin and the like) and may also be used.

Promoter regulatory elements may also be active during a certain stage of the plants' development as well as active in specific plant tissues and organs. Examples of such include but are not limited to pollen-specific, embryo specific, corn silk specific, cotton fiber specific, root specific, seed endosperm specific promoter regulatory elements and the like. Under certain circumstances it may be desirable to use an inducible promoter regulatory element responsible for expression of genes in response to a specific signal, such as: physical stimulus (heat shock genes); light (RUBP carboxylase); hormone (Em); metabolites; and stress. Other desirable transcription and translation elements functional in plants may also be used. Numerous plant-specific gene transfer vectors are known and available to the skilled artisan.

Heterologous DNA can be introduced into regenerable plant cell cultures via whiskers-mediated transformation. While a general description of the process can be found in US Patents 5,302,523 and 5,464,765, both to Zeneca, no protocols have been published to date for whisker-mediated transformation of Type I, Type II, hypocotyl or cotyledon regenerable plant cell lines.

In whisker-mediated transformation, DNA uptake into plant material is facilitated by very small, elongated, needle-like particles comprised of a biologically inert material. When said particles are agitated in the presence of DNA and plant cell lines, one or more of the particles produce small punctures in the regenerable plant cell aggregates thereby allowing said aggregates to uptake the DNA. Cells which have taken up the DNA are considered

to be transformed. Some transformed cells stably retain the introduced DNA and express it.

The elongated needle-like particles used in plant cell transformation are termed "whiskers" and are preferably made of a high density material such as silicon carbide or silicon nitride; however, any material having a needle-like structure wherein the size of said structure is smaller than the cell intended to be transformed is within the scope of the invention. More preferably, whiskers are made of silicon carbide and are either Silar SC-9 or Alfa Aesar as described herein.

Before transformation, plant cell lines are preferably placed onto an osmotic medium and allowed to incubate thereby enhancing transformation over non-osmotically treated cells. Most preferred is an osmotic medium having therein about 36.4 g/L sorbitol and about 36.4 g/L mannitol. These tissue are maintained on said medium for about 4 h before whisker-mediated transformation. Incubation on medium with osmoticum after transformation at the discretion of the artisan. However, transformation of non-osmotically treated plant cells aggregates and tissues is also within the scope of this invention.

Callus cultures used herein for generation of transgenic plants should generally be about 3 weeks to about 20 weeks old depending on the culture type. Typically, callus should be about midway between transfer periods and thus beyond any lag phase that might be associated with transfer to a new media or before reaching any stationary phase typically associated with a culture being on a plate for an extended period of time. However, plant material can be taken before or after sub-culturing; therefore, harvest timing is not generally believed to be

critical to practicing the invention as disclosed herein.
The amount of callus tissue used in each transformation
can vary with amounts of about 100 mg to about 500 mg
preferred, about 100 mg to about 250 mg being more
5 preferred and about 200 mg tissue being most preferred.

For transformation, whiskers are typically placed in
a small container, such as a conical or microfuge tube and
the like, wherein is placed a mixture comprising the DNA
construct of interest, a liquid medium, and callus tissue.
10 The order in which materials are added is not significant
to practicing the invention as disclosed herein.
Thereafter, the container is sealed and agitated. Unlike
particles used in biolistic transformation of plant tissue
(Sanford et al., 1990 Physiol. Plantarum, 79:206-209; and
15 US Patent 5,100,712), whiskers do not require any special
pretreatment with DNA carriers or precipitants prior to
use such as CaCl_2 , spermidine, sheared salmon sperm DNA and
the like.

Agitation time used in the transformation process can
20 vary and is typically from between about 10 sec to about
160 sec. The amount of whiskers added per transformation
can also vary from between about 1 mg to about 4 mg per
tube. An inverse relationship is observed between the
amount of whiskers added and the agitation time needed to
25 obtain optimal transformation. Therefore, the amount of
whiskers added and the agitation time needed to achieve
transformation is determinable by one having skill in the
art. In addition, the volume of liquid medium added can
vary from about 200 μL to about 1000 μL , with about 200 μL
30 being preferred. Moreover, the amount of heterologous DNA
added can vary from a preferred amount of about 10 μL to
about 100 μL of 1 mg/mL solution. The volume of DNA added
is not as critical of factor to the invention as disclosed

herein as the final DNA concentration. However, preferred final DNA concentrations are from about 0.03 $\mu\text{g}/\mu\text{L}$ to about 0.14 $\mu\text{g}/\mu\text{L}$. The scope of the present invention is not intended to be limited to said container size, the amount or concentration of heterologous DNA added, the volume of heterologous DNA added, the amount of the liquid medium added, the amount of callus material added or the amount of whiskers added as disclosed herein. The scope of the invention is also not intended to be limited by the instrumentation used to agitate the mixture or whether agitation is accomplished by manual or mechanical means.

Once the plant cell lines have been perforated and the heterologous DNA has entered therein, it is necessary to identify, propagate, and select those cells which not only contain the heterologous DNA of interest but are also capable of regeneration. Said cells and plants regenerated therefrom can be screened for the presence or absence of the heterologous DNA by various standard methods including but not limited to assessment of reporter gene expression. Alternatively, transmission of a selectable marker gene along with or as part of the heterologous DNA allows those cells containing said DNA to be identified by use of a selective agent.

Selection of only those cells containing and expressing the heterologous DNA of interest is a critical step in production of fertile, transgenic plants. Selection conditions must be chosen in such a manner as to allow growth of transformed cells while inhibiting growth of untransformed cells, which initially, are far more abundant. In addition, selection conditions must not be so severe as to cause transformed cells to lose their plant regenerability, future viability or fertility. A skilled artisan can easily determine appropriate

conditions for selecting transformed cells expressing a particular selectable marker by performing growth inhibition curves. Growth inhibition curves are generated by plotting cell growth versus selective agent concentration. Typically, selective agent concentrations are set at a concentration whereby almost all non-transformed cells are growth inhibited but yet are not killed. Preferred are selective agent concentrations wherein 90-99% of non-transformed cells are growth inhibited but yet not killed. Most preferred are selective agent concentrations wherein 97-99% of non-transformed cells are growth inhibited but yet not killed.

Transformed callus tissues transferred and exposed to selective agents are generally incubated on solid medium supportive of growth. The medium preferred for each type of tissue has been well defined in the art. After initial exposure to selective agents, tissues are transferred periodically to fresh medium while maintaining selective agent concentrations. After transformed cell mass has essentially doubled in size, masses showing the most growth and appearing to be healthy are selected and transferred to fresh medium having selective agent concentrations wherein non-transformed cells will be killed. Repeated selection and transference of growing cells to fresh medium result eventually in a selected group of cells comprised almost exclusively of transformed cells containing the heterologous DNA of interest.

Regeneration, while important to the present invention, may be performed in any conventional manner available to the skilled artisan. If cells have been transformed with selectable marker gene, the selective agent may be incorporated into the regeneration media to further confirm that the regenerated plantlets are

transformed. After subsequent weeks of culturing, regenerated plantlet immune to the selective agent can be transferred to soil and grown to maturity.

Callus and plant derived therefrom can be identified
5 as transformants by phenotypic and/or genotypic analysis. For example, if an enzyme or protein is encoded by the heterologous DNA, enzymatic or immunological assays specific for the particular enzyme or protein can be used. Other gene products may be assayed by using suitable
10 bioassays or chemical assays. Other techniques include analyzing the genomic component of the plant using methods as described by Southern ((1975) J. Mol. Biol., 98:503-517), polymerase chain reaction (PCR) and the like.

Plants regenerated from transformed callus are
15 referred to as the R0 generation or R0 plants. Seed produced by various sexual crosses from plants of this generation are referred to as R1 progeny. R1 seed are then germinated to produce R1 plants. Successful transmission and inheritance of heterologous DNA to R1
20 plants and beyond should be confirmed using the methods described herein.

Generally, the commercial value of transformed corn and progeny thereof will be of greatest value if the heterologous DNA can be incorporated into many different
25 hybrids. This may be achieved by incorporating the heterologous DNA into a large number of parental lines directly as described herein by creating plant cell aggregates of said lines and transforming said lines with whiskers. In addition, this may also be accomplished by
30 crossing initial transgenic fertile plants to normal elite inbred lines and then crossing the progeny thereof back to the normal parent. Progeny from this cross will segregate such that some plants will contain the heterologous DNA of

5

10

15

ESTABLISHMENT OF TYPE I and TYPE II MAIZE CALLUS TISSUE

20

cultures were used for transformation experiments described herein.

For Type II callus initiation, plants of a hybrid made by inter-mating two S₁ lines derived from a B73 x A188 cross (Armstrong et al., 1991, Maize Genet. Coop. News Lett., 65:92-93) were grown under standard greenhouse conditions and self- or sib-pollinated (Petolino and Genovesi, 1994, The Maize Handbook, pp. 701-704). Ten to 14 days after pollination (DAP), developing ears were removed and surface sterilized with 70% (v/v) ethanol for 2 min followed by soaking in 20% (v/v) commercial bleach for 0.5 h containing a few drops of LIQUI-NOX, then rinsed several times with sterile H₂O. Immature embryos (1.5 - 3.0 mm) were then isolated and placed onto initiation medium [N6 basal salts and vitamins (Chu, 1978, Proc. Symp. Plant Tissue Culture, pp. 43-56), 20 g/L sucrose, 2.9 g/L L-proline, 100 mg/L enzymatic casein hydrolysate (ECH), 37 mg/L Fe-EDTA, 10 mg/L AgNO₃, 1 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), and 2.5 g/L GELRITE (Schweizerhall, South Plainfield, NJ) at pH 5.8]. After 2-3 weeks in the dark at 28° C, Type II callus with numerous globular and elongated somatic embryos was obtained. An additional 2-3 selective subcultures were performed to enrich Type II cultures with desirable morphology as described in Welter et al., 1995, Plant Cell Rep. 14:725-729. Once Type II morphology was established (4-6 weeks), callus was transferred to maintenance medium (initiation medium containing 6 mM L-proline and no AgNO₃). Type II callus was used for transformation after about 16-20 weeks from culture initiation.

EXAMPLE 2

CONSTRUCTION OF THE PLASMID pUGN81-3 AND pDAB418

The maize expression vector, pUGN81-3, containing the ubiquitin promoter regulatory element driving the β -glucuronidase gene was used as disclosed herein. Plasmid pUGN81-3 was a 8730 base pairs double stranded plant transformation vector composed of the following sequences in clockwise order. Nucleotides 1 to 17 encoded a polylinker having the sequence AGCTTCCGCG GCTGCAG.

5 Nucleotides 18 to 2003 of pUGN81-3 were the maize ubiquitin promoter and first intron thereof and were PCR amplified from genomic DNA of maize genotype B73 (Christensen et al., (1992) Plant Mol. Biol. 18:675-689). Nucleotides 2004 to 2022 of pUGN81-3 encoded a polylinker

15 having the sequence GGTACCCCCG GGGTCGACC. Nucleotides 2023 to 4154 of pUGN81-3 corresponded to nucleotides 2551 to 4682 of plasmid pBI101 (Clontech, Palo Alto, CA) followed by a polylinker having the sequence ATCGGGAATT AAGCTTGCAT GCCTGCAGGC CGGCCTTAAT TAA which corresponded to

20 bases 4155 to 4197 of pUGN81-3. Nucleotides 4198 to 4264 of pUGN81-3 corresponded to: GCGGCCGCTT TAACGCCCGG GCATTTAAAT GGCGCGCCGC GATCGCTTGC AGATCTGCAT GGGTG. Nucleotides 4265-4776 of pUGN81-3 comprised the double-enhanced 35S promoter, with nucleotides 4265 to 4516

25 corresponding to nucleotides 7093 to 7344 of the Cauliflower Mosaic Virus genome (Franck et al., (1980) Cell 21:285-294). Nucleotides 4525 to 4776 of pUGN81-3 were a duplication of nucleotides 4265 to 4516 with the linker CATCGATG comprising nucleotides 4517 to 4524

30 between the duplicated sequence. Nucleotides 4777 to 4871 of pUGN81-3 corresponded to bases 7345 through 7439 of the Cauliflower Mosaic Virus genome (Franck et al., (1980) Cell 21:285-294). Nucleotides 4872 to 4891 comprised the

linker GGGGACTCTA GAGGATCCAG. Nucleotides 4892 to 5001 of pUGN81-3 corresponded to nucleotides 167 to 277 of the Maize Streak Virus genome with base 187 absent (Mullineaux et al., (1984) EMBO J. 3:3063-3068). Nucleotides 5002 to 5223 corresponded to the modified first intron of the maize alcohol dehydrogenase gene (Adh1-S) (Dennis et al., (1984) Nucleic Acids Res. 12:3983-4000). The modification resulted in removal of 343 nucleotides (bases 1313 to 1656) with bases 1222 to 1312 (intron 5' end) and nucleotides 1657 to 1775 (intron 3' end) of the maize Adh1-S gene remaining. Nucleotides 5224 to 5257 of pUGN81-3 corresponded to Maize Streak Virus (MSV) nucleotides 279 to 312. Both sections of the Maize Streak Virus, hereinafter MSV, sequence comprised the untranslated leader of the MSV coat protein V2 gene, and were interrupted in plasmid pUGN81-3 by the modified Adh1 intron. Nucleotides 5258 to 5814 of plasmid pUGN81-3 corresponded to nucleotides 29 to 585 of the phosphinotricin acetyl transferase (BAR) gene of *Streptomyces hygroscopicus* (White et al., (1989) Nucleic Acids Res. 18:1062). To facilitate cloning, nucleotides 34 and 575 of the published sequence were changed from A and G to G and A, respectively. This sequence served as the selectable marker. Nucleotides 5815 to 5819 comprised the linker GATCT. Nucleotides 5820 to 6089 of pUGN81-3 corresponded to nucleotides 4414 to 4683 of plasmid pBI101 (Clontech, Palo Alto, CA) followed by the linker sequence ATCGG. The remaining sequence of pUGN81-3 (nucleotides 6095 to 8730) corresponded to the reverse complement of pUC19 (Yanish-Perron et al., (1985) Gene 33:103-119).

The maize expression vector, pDAB418, contained the ubiquitin promoter regulatory element driving the β -glucuronidase gene was used some expression studies. In

addition this plasmid carried a second gene which served as a plant selectable marker. Plasmid pDAB418 was a 10,149 base pairs double stranded plant transformation vector composed of the following sequences in clockwise order. Nucleotides 1 to 31 had the nucleotide sequence AATTCATCGA AGCGGCCGCA AGCTTCCGCG G. Nucleotides 32 to 2023 of pDAB418 were the maize ubiquitin (Ubi1) promoter and first intron, and were PCR amplified from genomic DNA of maize genotype B73 (Christensen et al., (1992) Plant Mol. Biol. 18:675-689). Nucleotides 2024 to 2042 of pDAB418 comprised the linker sequence GGTACCCCCCG GGGTCGACC. Nucleotides 2043 to 3894 of pDAB418 corresponded to nucleotides 2551 to 4402 of plasmid pBI101 (Clontech, Palo Alto, CA) followed by the linker sequence TGGGGAATTG (bases 3895 to 3904). Nucleotides 3905 to 4174 OF pDAB418 corresponded to 4414 to 4683 of pBI101. Nucleotides 4175 to 4192 were composed of the linker sequence ATCGGGAATT AAGCTTGG. Base 4193 through 6184 was composed of a second copy of the maize ubiquitin promoter and first intron as describe above. This sequence was followed by the linker sequence GTCGGGGATC TA (6185 to 6196). Nucleotides 6197 to 6753 of plasmid pDAB418 corresponded to nucleotides 29 to 585 of the phosphinotricin acetyl transferase (BAR) gene of *Streptomyces hygroscopicus* (White et al., (1989) Nucleic Acids Res. 18:1062). To facilitate cloning, nucleotides 34 and 575 of the published sequence were changed from A and G to G to A, respectively. This sequence served as the selectable marker and was regulated by the maize ubiquitin promoter. Nucleotides 6754 to 6758 were composed of the linker TAACC. Nucleotides 6759 through 7472 functioned as the 3' polyadenylation sequence and includes bases 21728 through 22441 from the *Agrobacterium*.

tumefaciens octopine Ti plasmid pTi15955 (Barker et al. (1983) Plant Mol. Biol. 2, 335-350. Sequence 7473 through 7504 was composed of the polylinker GGAATTCATC GATATCTAGA TGTCGAGCTC GG. The remaining sequence of pDAB418

5 (nucleotides 7505 to 10149) corresponded to the reverse complement of nucleotides from the plasmid backbone derived from pUC19 (Yanish-Perron et al., (1985) Gene 33:103-119).

EXAMPLE 3

10 WHISKERS PREPARATION, OPTIMIZATION, AND TYPE I CORN CALLUS TRANSFORMATION

Type I callus was produced from different inbred lines. For testing, about 250 mg samples of each Type 1 callus was placed into 2.0 mL microfuge tubes (Brinkman
15 Instuments, Inc., Westbury, NY), 300 mL of Type I medium consisting of N6 macro-nutrients and vitamins (Chu, 1978, Proc. Symp. Plant Tissue Culture, pp. 43-56), B5 micro-nutrients (Gamborg et al., Exp. Ce.. Res. 50:151-158), 30 g/L sucrose, 2.9 g/L L-proline, 100 mg/L myo-inositol, 100
20 mg/L casein hydrolysate, 37 mg/L NaFeEDTA, 8 mg/L dicamba, 0.8 mg/L 2,4-D, 4.9 mg/L AgNO₃], 20 µL of DNA solution [1.0 mg/mL pDAB418 in 10 mM Tris, 1 mM EDTA, pH 8.0], and 200 µL of a 40 µg/µL (SC-9) whisker suspension was added. Each tube was then agitated for 20 sec using a Vortex-
25 Genie 2™ and the callus was transferred back to Type I medium solidified with 2.5 g/L GELRITE and allowed to incubate for 16 h. Afterwards, callus was assayed for GUS expression as described, *infra.*, with results summarized in Table 1.

30 TABLE 1. GUS expression in Type I callus.

Genotype	GUS Expression Units per Sample
H99	31 ± 26
Pa91	22 ± 8
139/39	3 ± 3

Transient GUS expression in Type I callus following whisker treatment was observed in all genotypes tested.

EXAMPLE 4

WHISKERS PREPARATION, OPTIMIZATION, AND TYPE II CORN CALLUS TRANSFORMATION

5 A sterilized suspension of silicon carbide whiskers was prepared by taking about 40 mg of dry whiskers (Silar SC-9 from Advanced Composite Materials Corp., Greer, SC; Alfa Aesar from Johnson-Matthey, Ward Hill, MA) and
10 placing them into a pre-weighed 2.0 mL polypropylene tube. The tube was then re-weighed to determine the amount of whiskers added, followed by autoclaving. Immediately before use, a 40 mg/mL whisker suspension was made by adding maintenance medium, *supra.*, and vortexing at high
15 speed for 1 min.

In the transformation experiments, either 200 or 500 mg samples of Type II callus from different immature embryo-derived lines were placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ).
20 Into each was added 500 μ L of liquid maintenance medium [minus GELRITE], 80 μ L of DNA solution [0.5 μ g/ μ L pDAB418 in 10 mM Tris, 1 mM EDTA, pH 8.0], and 50 μ L of a 40 mg/mL (SC-9) whisker suspension. Each tube was then agitated for 20 sec using a Vortex-Genie 2[™] (Scientific Industries,
25 Bohemia, NY). Negative control experiments were also performed. In one case, callus tissue was treated with whiskers in the absence of DNA. In the other cases, callus tissue was treated with DNA in the absence of whiskers. Regardless of the treatment, callus was then
30 transferred back to solid maintenance medium having 2.5 g/L GELRITE and allowed to incubate for 16 h in the dark at 28° C. Afterwards, callus was placed into GUS assay solution [0.2 M sodium phosphate pH 8.0, 0.1 mM each of

potassium ferricyanide and potassium ferrocyanide, 1.0 M sodium EDTA, 0.5 mg/mL 5-bromo-4-chloro-3-indolyl- β -D-glucuronide, and 0.6% v/v Triton x-100]. GUS expression, defined as counts of blue sectors per sample, was measured after 48 h of incubation at 37° C in the dark and is summarized in Table 2.

TABLE 2. GUS expression following whisker-mediated transformation of callus at a fixed DNA concentration (0.0635 μ g/ μ L Final concentration)

Callus Amount/tube Sample	GUS Expression Units Per
200 mg callus + whiskers + DNA	79 \pm 22
200 mg callus + whiskers - DNA	0
200 mg callus - whiskers + DNA	0
500 mg callus + whiskers + DNA	51 \pm 20
500 mg callus + whiskers - DNA	0
500 mg callus - whiskers + DNA	0

GUS expression was only observed when whiskers and DNA were both agitated in the presence of callus. Slightly higher transient GUS expression was observed when 200 mg of callus per tube was used as compared to 500 mg per tube.

To test the effect of different DNA concentrations, about 200 mg samples of Type II callus from different immature embryo-derived lines were placed into a 17 x 100 mm culture tube (Falcon 2059, Becton Dickinson, Lincoln Park, NJ). Into each was added 250 μ L of liquid maintenance medium, either 10, 25, or 50 μ L of DNA (1.0 μ g/ μ L pUGN81-3 in 10 mM Tris, 1 mM EDTA, pH 8.0), and 50 μ L of a 40 mg/mL (SC-9) whisker suspension. Each tube was then agitated for 20 sec using a Vortex-Genie 2™ (Scientific Industries, Bohemia, NY). The callus was then transferred back to semi-solid maintenance medium and allowed to incubate for 16 h in the dark at 28° C.

Afterwards, callus was assayed for GUS expression as previously described with the results shown in Table 3.

TABLE 3. GUS expression following whisker-mediated transformation of 200 mg callus at varied DNA concentrations.

Final DNA Concentration Sample	GUS Expression Units Per
0.032 $\mu\text{g}/\mu\text{L}$	105 \pm 73
0.077 $\mu\text{g}/\mu\text{L}$	86 \pm 57
0.143 $\mu\text{g}/\mu\text{L}$	96 \pm 76

As little as 10 μg of DNA per tube (0.032 $\mu\text{g}/\mu\text{L}$) containing about 200 mg of callus resulted in transient GUS expression. Increasing the amount of DNA did not appear to significantly increase transient GUS expression.

About 500 mg samples of Type II callus representing different immature embryo-derived lines were prepared as described herein and placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ). Into each was added 500 μL of liquid maintenance medium and 50 μL of a DNA solution [0.5 $\mu\text{g}/\mu\text{L}$ pDAB418 in 10 mM Tris, 1 mM EDTA, pH 8.0]. The callus/DNA mixture was then allowed to incubate for 1 h either at room temperature or on ice. Callus was incubated in the absence of DNA as a negative control. Afterwards, 50 μL of a 40 $\mu\text{g}/\mu\text{L}$ (SC-9) whisker suspension was added and vortexed as described herein. The callus was then transferred back to maintenance medium solidified with 2.5 g/L GELRITE and allowed to incubate for 16 h in the dark at 28° C. Afterwards, the callus was assayed for GUS expression as described previously. Results are summarized in Table 4.

TABLE 4. GUS expression following whisker-mediated transformation of callus pre-incubated with DNA for 1 h.

DNA Incubation Sample	GUS Expression Units Per
DNA + callus + whiskers	70 ± 65
DNA + callus → 1 h(ice) → whiskers	23 ± 37
DNA + callus → 1 h(RT) → whiskers	60 ± 27

Incubation of callus with DNA before whisker agitation resulted in lower transient GUS expression, especially when the incubation was at room temperature. Best results were observed when DNA was added immediately prior to whisker treatment.

About 200 mg samples of Type II callus representing different immature embryo-derived lines were either placed onto osmotic medium [maintenance medium containing 36.4 g/L sorbitol and 36.4 g/L mannitol] and allowed to incubate for 4 h or maintained on maintenance medium. The callus was then placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) and into each was added 200 µL of either liquid osmotic medium or liquid maintenance medium, 20 µL of DNA solution [1.0 µg/µL pUGN81-3 in 10 mM Tris, 1 mM EDTA, pH 8.0], and 50 µL of 40 µg/µL (Alfa) whisker suspension. Each tube was then agitated for 20 sec using a Vortex-Genie 2™, callus was transferred back to either osmotic medium or maintenance medium solidified with 2.5 g/L GELRITE and allowed to incubate for 16 h in the dark at 28° C. Callus was then tested for GUS expression as described previously. Results are summarized in Table 5.

TABLE 5. GUS expression of callus tissue having been osmotically treated before and/or after whisker-mediated transformation.

Osmotic Treatment	GUS Expression Units Per Sample
None	72 ± 19
Before Only	245 ± 80
After Only	67 ± 37
Before and After	222 ± 83

Transient GUS expression was highest in those cultures which included an osmotic treatment before whisker agitation.

About 200 mg samples of Type II callus representing different cell lines were placed onto osmotic medium and allowed to incubate for 4 h. The callus was then placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) and into each was added 200 µL of liquid osmotic medium [minus GELRITE], 20 µL of DNA solution [1.0 mg/mL pUGN81-3 in 10 mM Tris, 1 mM EDTA, pH 8.0] and either 25, 50, or 100 µL of a 40 µg/µl (Alfa) whisker suspension. Each tube was then agitated for either 20- or 60 sec using a Vortex-Genie 2™. The callus was then transferred back to liquid osmotic medium, allowed to incubate for 16 h in the dark at 28° C, and then assayed for GUS expression as described previously. Results are summarized in Table 6.

TABLE 6. GUS expression of callus tissue transformed with various amounts of whiskers for 20- and 60 sec.

Whisker Sample Amount seconds	GUS Expression Units Per	
	20 seconds	60
25 µL (1 mg/tube)	61 ± 36	136 ±
50 µL (2 mg/tube)	70 ± 46	183 ±
100 µL (4 mg/tube)	155 ± 92	175 ±

Averaged over all whisker amounts, highest GUS expression was observed following 60 sec agitation compared to 20 sec. However, a 60 sec agitation appeared to substantially damage the callus. High GUS expression was also observed when 100 μ L of the whisker suspension was used (4 mg whiskers/tube). Our data indicate that increased whisker amounts can compensate for agitation time and visa versa. The most preferred conditions was 100 μ L of whiskers vortexed for 20 sec.

About 500 mg samples of Type II callus representing different cell lines were placed onto osmotic medium and allowed to incubate for 4 h. The callus was then placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) and into each was added 1000 μ L of liquid osmotic medium, 27.7 μ L of DNA solution [0.72 μ g/ μ L pUGN81-3 in 10 mM Tris, 1 mM EDTA, pH 8.0], and 200 μ L of 40 μ g/ μ L whisker suspension using either Silar SC-9 (Advanced Composite, Greer, SC) or Alfa Aesar (Johnson Matthey, Ward Hill, MA) whiskers. Each tube was then agitated for 20 sec using a Vortex-Genie 2TM and the callus was transferred back to osmotic medium solidified with 2.5 g/L GELRITE and allowed to incubate for 16 h at 28° C. Afterwards, callus was assayed for GUS expression as described previously with results summarized in Table 7.

TABLE 7. Comparison of whisker type on transformation.

Whisker Type Sample	GUS Expression Units per
Silar SC-9	113 \pm 48
Alfa Aesar	51 \pm 9

Although both worked, better transient GUS expression was observed using Silar SC-9 whiskers than Alfa Aesar whiskers.

About 500 mg samples of Type II callus representing different cell lines were placed onto osmotic medium and allowed to incubate for 4 h. The callus was then placed into either 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) or 15 mL conical centrifuge tubes (Corning 25319-15, Corning, NY). Into each was added 1000 μ L of liquid osmotic medium, 20 μ L of DNA solution [1.0 mg/mL pUGN81-3 in 10 mM Tris, 1 mM EDTA, pH 8.0], and 200 μ L of 40 μ g/ μ L (Alfa) whisker suspension. Each tube was then agitated for 20 sec using either a Vortex-Genie or a Caulk Vari-Mix II (Estrada Dental, Rancho Cucamonga, CA). Callus was then transferred back to solid osmotic medium, allowed to incubate for 16 h and then assayed for GUS activity (Table 8) as described previously. High levels of transient GUS expression were observed following agitation using both the Vortex and Vari-Mix; however, vortexing appeared to be somewhat better.

TABLE 8. The effect of vessel and agitator type on the transformation of callus tissue with whiskers.

Vessel Type	GUS Expression Units Per Sample	
	Vortex	Vari-Mix
Falcon	418 \pm 202	256 \pm 261
Corning	395 \pm 362	353 \pm 271
Agitator Average	407 \pm 294	304 \pm 270

EXAMPLE 5

PRODUCTION AND REGENERATION OF STABLY TRANSFORMED TRANSGENIC MAIZE PLANTS

About 200 mg samples of Type II callus were placed onto osmotic medium and allowed to incubate for 4 h. The

callus was then placed into 17x100 mm culture tubes (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) and into each was added 200 μ L of liquid osmotic medium, 20 μ L of DNA solution [1.0 μ g/ μ L pUGN81-3 in 10 mM Tris, 1 mM EDTA, 5 pH 8.0], and 100 μ L of 40 μ g/ μ L (Alfa) whisker suspension. Each tube was then agitated for 20 sec using a Vortex-Genie 2TM, the callus was transferred back to osmotic medium solidified with 2.5 g/L GELRITE and allowed to recover for ca. 48 h prior to selection.

10 The whisker-treated callus was placed onto selection medium [maintenance medium with 30 mg/L BastaTM (Hoechst, Frankfurt, Germany) and no enzymatic casein hydrolysate or L-proline] and transferred to fresh selection medium every four weeks for about 3-4 months. After 10-20 weeks, 15 actively growing colonies were isolated and sub-cultured onto fresh selection medium every two weeks to bulk-up callus prior to regeneration.

For plant regeneration, callus was transferred to induction medium, *infra.*, and incubated at 28° C, 16h/8h 20 light/dark photoperiod in low light (13 mE/m²/sec) for one week followed by 28° C, 16h/8h light dark photoperiod in high light (40 mE/m²/sec) for one week provided by cool white fluorescent lamps. The induction medium was composed of MS salts and vitamins (Murashige and Skoog, 25 1962, *Physiol. Plant.* 15:473-497), 30 g/L sucrose, 100 mg/L myo-inositol, 5 mg/L benzyl amino purine, 0.025 mg/L 2,4-D, 2.5 g/L GELRITE and adjusted to pH 5.7. Following this two-week induction period, callus was transferred to regeneration medium and incubated in high light (40 30 mE/m²/sec) at 28° C. The regeneration medium was composed of MS salts and vitamins, 30 g/L sucrose, and 2.5 g/L GELRITE adjusted to pH 5.7. The callus was sub-cultured

to fresh regeneration medium about every two weeks until plantlets appeared. Both induction and regeneration medium contained 30 mg/L Basta™. Plantlets were transferred to 10 cm pots containing approximately 0.1 kg of dry Metro-Mix 360 (The Scotts Co, Marysville, OH), placed in a greenhouse, moistened thoroughly, and covered with clear plastic cups for 2-4 days. At the 3-5 leaf stage, plants were transplanted to 5-gallon pots containing soil and grown to maturity.

10

EXAMPLE 6

SOUTHERN ANALYSIS OF TRANSFORMED CALLUS AND PLANT TISSUES

BASTA resistant lines were characterized by Southern analysis to confirm the presence of the transgene using a DNA probe specific for the coding region of the gene of interest. DNA from both callus and leaf material was analyzed.

For callus, the material was soaked in distilled water for 30 min. and transferred to a new petri dish prior to lyophilization. Leaf material from plants was harvested at the 6-8 leaf stage. Genomic DNA was prepared from lyophilized tissue as described by Saghai-Maroo et al. ((1984) Proceed. Nat. Acad. Sci. USA 81:8014-8018). Eight µg of each DNA was digested with the restriction enzyme(s) specific for each plasmid construct using conditions suggested by the manufacturer (Bethesda Research, Gaithersburg, MD) and separated by electrophoresis on a 0.8% agarose gel. The DNA was then blotted onto nylon membranes as described by Southern ((1975) J. Mol. Biol., 98:503-517). A 1.9 kb DNA probe specific for the GUS coding region was prepared using an Oligolabelling Kit (Pharmacia LKB, Piscataway, NJ) with 50 mCi of [α-³²P]dCTP (Amersham Life Science, Arlington Heights, IL). The radioactive probe was then hybridized to

the genomic DNA on the blots in 45 mL of minimal hybridization buffer [10% polyethylene glycol, 7% sodium dodecyl sulfate (SDS), 0.6x SSC where 1X SSC is 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0, 10 mM sodium phosphate, 5 mM EDTA and 100 µg/mL denatured salmon sperm DNA] overnight at 60° C. After hybridization, blots were washed at 60° C in 0.25X SSC and 0.2% SDS for 45 min., blotted dry and exposed to XAR-5 film (Kodak, Rochester, NY) overnight on two intensifying screens (DuPont, Newark, DE).

Plants regenerated from transgenic callus lines were tested via Southern analysis and all were found to have the 4.2 kb hybridization product. The hybridizing product was expected to contain the ubiquitin-1 promoter/intron, the GUS coding sequence, and the nos 3' untranslated region. Individual plants regenerated from a given culture all displayed identical hybridization patterns.

EXAMPLE 7

ESTABLISHMENT OF EMBRYOGENIC TYPE I RICE CALLUS TISSUE

For initiation of rice embryogenic Type I callus cultures, mature seeds of a *Oryza sativa* L. cv. Japonica, Taipei 309, were dehusked and surface-sterilized in 70% (v/v) ethanol for 2-5 min followed by a 30-45 min soak in 50% (v/v) commercial bleach containing a few drops of LIQUI-NOX (Alconox, Inc., New York, NY). The seeds were then rinsed 3 times in sterile H₂O and placed on filter paper before being transferred to induction medium [N6 macro elements (Chu, 1978, Proc. Symp. Plant Tissue Culture, Peking Press, p 43-56), B5 micro elements and vitamins (Gamborg et al., 1968, Exp. Cell Res. 50:151-158), 300 mg/L casein hydrolysate, 500 mg/L L-proline, 500 mg/L L-glutamine, 30 g/L sucrose, 2 mg/L 2,4-D, and 2.5

g/L GELRITE, pH 5.8]. The seeds were cultured on induction medium and incubated in the dark at 28° C for 3 weeks. Afterwards, emerging primary callus induced from the scutellar region was transferred to fresh induction medium for further maintenance. In general, embryonic callus was selected based on its morphology after each subculture. The general morphology of rice Type I embryogenic callus emerged as hard, compact, nodular structures appearing, and in some cases not appearing, embryo-like. This material was termed Type I.

EXAMPLE 8

WHISKERS PREPARATION, OPTIMIZATION, AND RICE TYPE I CALLUS TRANSFORMATION

Immediately prior to DNA delivery, a sterile 50 µg/µL suspension of silicon carbide whiskers (Silar SC-9) was prepared in water as described previously. About 125-500 mg of Type I rice embryogenic callus being less than 4 months-old was selected and transferred into either a 15 mL Corning conical centrifuge tube or a 1.5 mL microfuge tube to which about 250 µL of liquid initiation medium (initiation medium without GELRITE) had been added. Then, about 10-160 µL of a 50 µg/µL whisker solution was added along with 25 µL of 1 µg/µL pDAB418 DNA solution immediately prior to vortexing for 15-120 sec with Vortex Genie 2™ at the highest speed possible. Afterwards, the callus was transferred to initiation medium with and without high osmoticum [0.2 M mannitol and 0.2 M sorbitol; Vain et al., 1993, Plant Cell Rep. 12:84-88] and incubated in the dark at 28° C for about 24 h before being examined for GUS activity.

Gus histochemical assays were conducted according to Jefferson et al., (1987, EMBO J 6: 3901-3907) and as described here. Tissues were placed in 24-well microtiter

plates (Corning, New York, NY) containing 500 μ L of assay buffer per well. The assay buffer consisted of 0.2 M sodium phosphate (pH 8.0), 0.1 mM potassium ferricyanide, 0.1 mM potassium ferrocyanide, 1.0 M sodium EDTA, 1.9 mM 5-bromo-4-chloro-3-indolyl- β -D-glucuronide, and 0.06% triton X-100. The plates were incubated in the dark for 1-2 days at 37° C before microscopically examining the tissue for expression. Silicon carbide whisker-mediated DNA delivery was measured in terms of GUS transient expression, i.e., GUS expression units (blue spots) per sample.

Many different transformation parameters were tested to determine the optimum conditions needed for transformation. Some of the parameters tested included: vortexing time, amount of the 50 μ g/ μ L whisker solution added, treatment with high osmoticum before and/or after whisker treatment, amount of tissue used per transformation, and vessel size. In all of these experiments, the protocols described herein were essentially the same unless otherwise mentioned.

The effect of vortexing time (15-60 sec) was studied for DNA delivery into rice embryogenic Type I callus. In this experiment, callus was treated with high osmoticum for about 4 h prior to transformation with whiskers and for about 16 h following whisker treatment. Callus was transformed by placing 250 mg of callus into a 15 mL Corning conical centrifuge tube, adding 1000 μ L of liquid initiation medium, 25 μ L of a 1 μ g/ μ L DNA solution (pDAB418), and 100 μ L of a 50 μ g/ μ L whisker solution. Multiple sample sets were vortexed for 15-, 30-, 60- and 120 sec and the results were quantitated as shown in Table 9.

Table 9. The effect of vortexing time on the transformation of rice Type I embryogenic callus.

Vortexing Time (Sec)	Gus Expression Units per Sample
15	113.5
30	107.5
60	90.5
120	72.5

As can be seen, DNA delivery and callus transformation was possible when vortexing for as little as 15 sec despite the hard and compact morphology of rice Type I embryogenic callus.

The effect of varying the amount of whiskers added was examined while maintaining a constant volume (1000 μ L) of liquid initiation medium. Callus samples were treated with high osmoticum before and after whisker-mediated transformation. About 250 mg of callus tissue was selected and transferred to a 15 mL Corning conical centrifuge tube to which 1,000 μ L of liquid initiation medium lacking GELRITE was added. Different amounts of a 50 μ g/ μ L whisker solution (10, 20, 40, 80, and 160 μ L) along with 25 μ L of a 1 μ g/ μ L DNA solution (pDAB418) were added prior to vortexing with Vortex Genie 2TM for 60 sec. Samples were then transferred back onto the initiation medium with high osmoticum overnight before histochemical GUS assays were performed. The results are described in Table 10. GUS expression was found to correlate with an increasing abundance of whiskers added.

Table 10. The effect of increasing the amount of whiskers while maintaining a constant volume of initiation medium.

Whisker Solution Added (μ L)	Gus Expression Units/Sample
10	2
20	1
40	9
80	6
160	6.5

The effect of high osmoticum on transformation efficiency was studied with the results summarized in Table 11. In this experiment, callus tissue was subjected to either no high osmoticum treatment or high osmoticum treatment after whisker-mediated transformation. About 250 mg of callus tissue was transferred to 1.5 mL microfuge tube. About 250 μ L of liquid initiation medium was added followed by 125 μ L of 50 μ g/ μ L whisker solution and 25 μ L of a 1 μ g/ μ L DNA solution (pDAB418). The samples were vortexed for 60 sec and transferred back onto the initiation medium overnight with or without high osmoticum before GUS assays were conducted.

Table 11. The effect of osmoticum treatment after vortexing and vortexing time on transformation and transgene expression of rice Type I embryogenic callus tissue.

Osmoticum After Transformation	mg Tissue	Vortex Sec.	GUS Expression Units Per Sample
+	250	30	48
+	250	60	44
+	500	30	37
+	500	60	44
-	250	30	87
-	250	60	107
-	500	30	96
-	500	60	120

Callus tissue with no high osmoticum treatment after whisker transformation resulted in higher expression levels compared to that where high osmoticum treatment occurred.

The amount of callus tissue added and vessel size (1.5 mL microfuge tube vs. 2.0 mL microfuge tube) used were studied. In this experiment, 125 or 250 mg of callus tissue were used while maintaining a constant volume of liquid initiation medium, i.e., 250 μ L. In all cases, 125 μ L of a 50 μ g/ μ L whisker solution along with 25 μ L of a 1 μ g/ μ L DNA solution (pDAB418) were added prior to vortexing with Vortex Genie 2™ for 60 sec. Following transformation, callus samples were incubated on the initiation medium with high osmoticum overnight before GUS assays were conducted. The results are summarized in Table 12.

Table 12. The effect of vessel size and the amount of tissue added on the whiskers-mediated transformation of rice Type 1 callus tissue.

mg Tissue	Vessel Size (mL)	Gus Expression Units/Sample
125	1.5	8
125	2.0	6
250	1.5	24
250	2.0	9.5

The use of 1.5 mL microfuge tube and 250 mg callus tissue per tube resulted in highest transient expression.

EXAMPLE 9

PLASMID DESCRIPTION FOR pUbiHyg

The rice expression vector, pUbiHyg, contained the maize ubiquitin promoter and first intron from the ubiquitin gene (Ubi1) regulatory element driving the hygromycin B phosphotransferase (resistance) gene as described by Gritz and Davies, (1983) Gene 25:179-188. Plasmid pUbiHyg was a 5991 base pairs double stranded plant transformation vector composed of the following sequences in clockwise order. Nucleotides 1 to 42 had the

sequence AAGCTTGCAT GCCTGCAGAT CTGCGGCCGC AAGCTTCCGC GG.
Nucleotides 43 through 2034 of pUbiHyg were the maize
ubiquitin promoter and first intron, and were PCR
amplified from genomic DNA of maize genotype B73
5 (Christensen et al., (1992) Plant Mol. Biol. 18:675-689)
Nucleotides 2035 to 2052 of pUbiHyg had the sequence
GGTACCCCCG GGTAGACC. Nucleotides 2053 through 3078 of
pUbiHyg corresponded to nucleotides 211 through 1236 of
the the hygromycin B phosphotransferase (resistance) gene
10 sequence (accession number K01193), with bases 2056 and
2057 of pUbiHyg modified from AA to GT to facilitate
future cloning. Bases 3079 through 3097 were composed of
the linker TAAGAGCTCG AATTTCCTCC. Bases 3098 through 3351
corresponded to nucleotides 4430 to 4683 of plasmid pBI101
15 (Clontech, Palo Alto, CA). The remaining sequence of
pUbiHyg (nucleotides 3352 to 5991) corresponded to
nucleotides from the plasmid backbone, (Yanish-Perron et
al., (1985) Gene 33:103-119).

EXAMPLE 10

20 PRODUCTION OF STABLY TRANSFORMED RICE TYPE I CALLUS TISSUE
AND REGENERATION OF TRANSGENIC PLANTS

For production of stable rice transgenic plants, rice
Type I callus tissue was produced as described herein and
subjected to whisker-mediated transformation. Typically,
25 250 mg of rice Type I callus, with or without high
osmoticum treatment for four hours, was added to a 1.5
microfuge tube. Also added were 250 μ L of liquid
initiation medium, 125 μ L of a 50 μ g/ μ L whisker solution,
and 25 μ L of a 1 μ g/ μ L DNA solution (pDAB 305 and UbiHyg
30 in 1:1 ratio). Vortexing was carried out for 60 sec as
described previously to ensure DNA delivery. Following
whisker-mediated DNA delivery, callus was transferred to
initiation medium, as described previously, with high

osmoticum for 24 h before transfer to selection medium. Selection medium consisted of initiation medium with 30 mg/L hygromycin (CalBiochem-Novabiochem Corporation, La Jolla, CA). After 2 weeks, cultures were transferred to
5 fresh selection medium having 50mg/L hygromycin (Li et al., 1993, Plant Cell Rep. 12: 250-255). Placing cultures on selection medium resulted in the formation of compact, white-yellow, embryogenic callus cultures after 30-45 days. These cultures were then transferred to pre-
10 regeneration (PR) medium having 50 mg/L hygromycin and maintained in the dark at 28° C. PR medium consisted of initiation medium with 2 mg/L benzyl aminopurine (BAP), 1 mg/L naphthalene acetic acid (NAA), and 5 mg/L abscisic acid (ABA). After 2 weeks, cultures were then transferred
15 to regeneration (RN) medium wherein RN medium was initiation medium with 3 mg/L BAP, and 0.5 mg/L NAA. Cultures on RN medium were incubated for at 28° C under high fluorescent lights (325-ft-candles) until plantlets emerged. When the plantlet shoots reached about 2 cm,
20 they were transferred to Magenta GA7 boxes (Magenta Corp., Chicago, IL) containing medium consisting of MS macro and micro nutrients along with B5 vitamins (Gamborg et al., 1968, Exp. Cell Res. 50:151-158) which had been diluted 1:1 with water, 10 g/L sucrose, 0.05 mg/L NAA, 50 mg/L
25 hygromycin, 2.5 g/L GELRITE, and adjusted to pH 5.8. When plantlets were established with well-developed root systems, they were transferred to soil/metromix 360 (1:1) and raised in the greenhouse (29°/24° C day/night cycle, 50-60% humidity, 12 h photoperiod) until maturity.
30 Southern analysis of these plants revealed the presence of the hygromycin gene indicating that they were indeed transformed. These plants were successfully grown to maturity in the greenhouse.

EXAMPLE 11

PLASMID DESCRIPTION FOR pSMGN179-3

The expression vector, pSMGN179-3, contained a modified derivative of the chimeric regulatory regions from the *Agrobacterium tumefaciens* opine synthase genes described by Gelvin and Hauptmann, (1995) Patent WO 95/14098 driving the β -glucuronidase gene. Plasmid pSMGN179-3 was a 5541 base pairs double stranded plant transformation vector composed of the following sequences in clockwise order: nucleotides 1 to 16 had the multiple cloning sequence from the plasmid backbone, pUC19 (Yanish-Perron et al., (1985) Gene 33:103-119) AATTCGAGCTCGGTAC. Nucleotides 17 to 57 of pSMGN179-3 were composed of linker fragment having the sequence CGGGCCCCCCCTCGAGGTCG ACGGTATCGA TAAGCTTGAT C. Bases 58 through 277 of pSMGN179-3 corresponded to the reverse complement of bases 13774 through 13993 of *Agrobacterium tumefaciens* Ti plasmid pTi15955 T-DNA (Barker et Gal. (1983) Plant Mol. Biol. 2:335-350). These bases corresponded to upstream activating sequences from the octopine synthase (ocs) gene. Bases 278 through 304 corresponded to linker DNA with the sequence CGAATTCCAA GCTTGGGCTG CAGGTCA. Bases 305 through 350 corresponded to the reverse complement of bases 21475 through 21520 of *Agrobacterium tumefaciens* Ti plasmid pTi15955 T-DNA (Barker et Gal. (1983) Plant Mol. Biol. 2:335-350). Bases 351 through 737 of pSMGN179-3 were composed of the reverse complement of bases 20128 through 20514 from *Agrobacterium* b-glucuronidase gene *tumefaciens* Ti plasmid pTi15955 T-DNA (Barker et Gal. (1983) Plant Mol. Biol. 2:335-350). The sequence of pSMGN179-3 from 305 through 737 included the Δ MAS promoter described by Gelvin and Haupman, (1995) Patent WO 95/14098. Bases 738 through 763 of pSMGN179-3 contain

linker sequence CTCTAGAACT AGTGGATCCG TCGACC. Bases 58 through 763 of pSMGN179-3 comprised the promoter and untranslated leader regulatory fusion as given in SEQ ID NO 1. Nucleotides 764 to 2615 of pSMGN179-3 corresponded to nucleotides 2551 to 4402 of plasmid pBI101 which encoded the β -glucuronidase gene (Clontech, Palo Alto, CA) (Jefferson et al., (1986) Plant Mol. Biol. Rep. 83(22):8447-8451). Bases 767 through 769 were modified from TTA to GTC. The β -glucuronidase gene was followed by the polylinker sequence TGGGGAATTG, corresponding to bases 2616 to 2625 of pSMGN179-3. Bases 2626 through 2895 corresponded to 4414 to 4683 of pBI101 which contained the sequence from the nopaline synthase 3' untranslated regions (Clontech, Palo Alto, CA) (Jefferson et al., (1986) Plant Mol. Biol. Rep. 83(22):8447-8451). The remaining sequence of pSMGN179-3 (nucleotides 4684 to 5541) corresponded the reverse complement of nucleotides from the plasmid backbone which was derived from pUC19 (Yanish-Perron et al., (1985) Gene 33:103-119).

20

EXAMPLE 12

TRANSFORMATION OF EMBRYOGENIC COTTON CALLUS

Approximately 250 mg of cotton callus from different seedling-derived lines were placed onto osmotic medium [maintenance medium plus 36.4 g/L sorbitol and 36.4 g/L mannitol] and incubated for 4 h in the dark. The callus was then placed into a 17 x 100 mm culture tube (Falcon 2059, Becton Dickinson, Lincoln Park, NJ) into which was added 250 μ L of liquid osmotic medium [minus GELRITE], 20 μ L of DNA (1.0 μ g/ μ L pSMGN179-3 in 10 mM Tris, 1 mM EDTA, pH 8.0), and 50 μ L of a 40 mg/mL (Alfa) whisker suspension. Each tube was then agitated for either 40, 80, or 160 sec using a Vortex-Genie 2TM (Scientific

Industries, Bohemia, NY). The callus was then transferred back to osmotic medium solidified with 2.5 g/L GELRITE and allowed to incubate at 28° C in the dark for 16 h.

Afterwards, it was assayed for GUS expression as

5 previously described with the results shown in Table 13.

TABLE 13. GUS expression following whisker-mediated transformation cotton callus using different agitation times.

10	Agitation Time Sample	GUS Expression Units Per
	40 sec	41 ± 21
	80 sec	59 ± 38
	160 sec	37 ± 28

15

GUS expression was observed in whisker-treated cotton callus. Increasing the agitation time from 40 to 160 sec did not appear to significantly increase expression.

EXAMPLE 13

20

PRODUCTION OF PLASMID pDAB 305

Plasmid pDAB305 was a 5800 bp plasmid that harbored a promoter containing a tandem copy of the Cauliflower Mosaic Virus 35S enhancer (35S), a deleted version of the *Adh1* intron 1, and the untranslated leader from the Maize Streak Mosaic Virus Coat Protein fused to the β -glucuronidase gene, which was then followed by the *nos* 3'UTR. It was made as follows:

The starting material was plasmid pUC13/35S(-343) as described by Odell et al. (1985 Nature 313:810-812). This plasmid comprised, starting at the 3' end of the *Sma* I site of pUC13 (Messing, 1983 in Methods in Enzymology, Wu, R. Ed. 101:20-78), and reading on the strand contiguous to the noncoding strand of the *lacZ* gene of pUC13, nucleotides 6495 to 6972 of CaMV, followed by the linker sequence CATCGATG (which encodes a *Cla* I recognition site), followed by CaMV nucleotides 7089 to 7443, followed

35

by the linker sequence CAAGCTTG, the latter sequence comprising the recognition sequence for Hind III, which was then followed by the remainder of the pUC13 plasmid DNA.

5 The plasmid pUC13/35S(-343) DNA was digested with Cla I and Nco I, the 3429 base pair (bp) large fragment was separated from the 66 bp small fragment by agarose gel electrophoresis, and then purified by standard methods. The DNA was digested with Cla I, and the protruding ends
10 were made flush by treatment with T4 DNA polymerase. The blunt-ended DNA was then ligated to synthetic oligonucleotide linkers having the sequence CCCATGGG, which included an Nco I recognition site. The ligation reaction was transformed into competent *E. coli* cells, and
15 a transformant was identified that contained a plasmid (named pOO#1) that had an Nco I site positioned at the former Cla I site. DNA of pOO#1 was digested with Nco I and the compatible ends of the large fragment were religated, resulting in the deletion of 70 bp from pOO#1,
20 to generate intermediate plasmid pOO#1 NcoE.

 The plasmid pOO#1 NcoE DNA was digested with EcoR V, and the blunt ends were ligated to Cla I linkers having the sequence CATCGATG. An *E. coli* transformant harboring a plasmid having a new Cla I site at the position of the
25 previous EcoR V site was identified, and the plasmid was named pOO#1 NcoE RV>Cla. This DNA was then digested with Cla I and Nco I, and the small (268 bp) fragment was purified from an agarose gel. This fragment was then ligated to the 3429 bp Cla I/Nco I fragment of pUC13/35S(-
30 343) prepared above, and an *E. coli* transformant that harbored a plasmid having Cla I/Nco I fragments 3429 and 268 bp was identified. This plasmid was named pUC13/35S En.

The plasmid pUC13/35S En DNA was digested with Nco I, and the protruding ends were made blunt by treatment with T4 DNA polymerase. The treated DNA was then cut with Sma I, and was ligated to Bgl II linkers having the sequence CAGATCTG. An *E. coli* transformant that harbored a plasmid in which the 416 bp Sma I/NcoI fragment had been replaced with at least two copies of the Bgl II linkers was identified, and named p35S En². The DNA structure of p35S En² was as follows: Beginning with the nucleotide that follows the third C residue of the Sma I site on the strand contiguous to the noncoding strand of the *lacZ* gene of pUC13; the linker sequence CAGATCTGCA GATCTGCATG GGCGATG, followed by CaMV nucleotides 7090 to 7344, followed by the Cla I linker sequence CATCGATG, followed by CaMV nucleotides 7089 to 7443, followed by the Hind III linker sequence CAAGCTT, followed by the rest of pUC13 sequence. This structure had the feature that the enhancer sequences of the CaMV 35S promoter, which laid in the region upstream of the EcoR V site in the viral genome (bases 7090 to 7344), had been duplicated. This promoter construct incorporated the native 35S transcription start site, which was 11 nucleotides upstream of the first A residue of the Hind III site.

For plasmids utilizing the 35S promoter and the *Agrobacterium* NOS Poly A sequences, the starting material for the first construct was plasmid pBI221, purchased from CLONTECH (Palo Alto, CA). This plasmid contained a slightly modified copy of the CaMV 35S promoter. Beginning at the 3' end of the Pst I site of pUC19 (Yanisch-Perron et al., 1985 Gene 33:103-119), and reading on the same strand as that which encodes the *lacZ* gene of pUC19, the sequence was comprised of the linker nucleotides GTCCCC, followed by CaMV nucleotides 6605 to

7439 followed by the linker sequence GGGGACTCTA GAGGATCCCC
GGGTGGTCAG TCCCTT. These bases were then followed by 1809
bp comprising the coding sequence of the *E. coli uidA*
gene, which encodes the β -glucuronidase (GUS) protein, and
5 55 bp of 3' flanking bases that are derived from the *E.*
coli genome (Jefferson, 1986 Proc. Natl. Acad. Sci.
83:8447-8451), followed by the Sac I linker sequence
GAGCTC, which was then followed by the linker sequence
GAATTTCCCC. These bases were followed by the RNA
10 transcription termination/polyadenylation signal sequences
derived from the *Agrobacterium tumefaciens* nopaline
synthase (NOS) gene, and comprised the 256 bp Sau3A I
fragment corresponding to nucleotides 1298 to 1554 of
DePicker et al. (1982 J. Molec. Appl. Genet. 1:561-573),
15 followed by two C residues, the Eco RI recognition
sequence GAATTC, and the rest of pUC19.

The plasmid pBI221 DNA was digested with EcoR I and
BamH I, and the 3507 bp fragment was purified from an
agarose gel. Plasmid pRAJ275 (CLONTECH) DNA was digested
20 with EcoR I and Sal I, and the 1862 bp fragment was
purified from an agarose gel. These two fragments were
mixed together, and complementary synthetic
oligonucleotides having the sequence GATCCGGATC CG and
TCGACGGATC CG were added. The fragments were ligated
25 together, and an *E. coli* transformant harboring a plasmid
having the appropriate DNA structure was identified by
restriction enzyme analysis. DNA of this plasmid, named
pKA881, was digested with Bal I and Eco RI, and the 4148
bp fragment was isolated from an agarose gel. DNA of
30 pBI221 was similarly digested, and the 1517 bp Eco RI/Bal
I fragment was gel purified and ligated to the above
pKA881 fragment, to generate plasmid pKA882. Then pKA882
DNA was digested with Sac I, the protruding ends were made

blunt by treatment with T4 DNA polymerase, and the fragment was ligated to synthetic BamH I linkers having the sequence CGGATCCG. An *E.coli* transformant that harbored a plasmid having BamH I fragments of 3784 and 1885 bp was identified and named pKA882B. Following, pKA882B DNA was digested with BamH I, and the mixture of fragments was ligated. An *E.coli* transformant that harbored a plasmid that generated a single 3783 bp fragment upon digestion with BamH I was identified and named p35S/NOS. This plasmid had the essential DNA structure of pBI221, except that the coding sequences of the GUS gene had been deleted. Therefore, CaMV nucleotides 6605 to 7439 were followed by the linker sequence GGGGACTCTA GAGGATCCCG AATTTCCTCC. The linker sequence was then followed by the NOS Polyadenylation sequences and the rest of pBI221.

The plasmid p35S/NOS DNA was digested with EcoR V and Pst I, and the 3037 bp fragment was purified and ligated to the 534 bp fragment obtained from digestion of p35S En² DNA with EcoR V and Pst I. An *E. coli* transformant was identified that harbored a plasmid that generated fragments of 3031 and 534 bp upon digestion with EcoR V and Pst I, and the plasmid was named p35S En²/NOS. This plasmid contained the duplicated 35S promoter enhancer region described for p35S En², the promoter sequences being separated from the NOS polyadenylation sequences by linker sequences that included unique Xba I and BamH I sites.

The MSV genomic sequence was published by Mullineaux et al., (1984 EMBO J. 3:3063-3068), and Howell (1984 Nucleic Acids Res. 12:7459-7375), and the transcript was described by Fenoll et al. (1988 EMBO J. 7:1589-1596).

The entire sequence, comprised 154 bp and was constructed in three stages by assembling blocks of synthetic oligonucleotides. First, complementary oligonucleotides having the sequence GATCCAGCTG AAGGCTCGAC AAGGCAGATC
5 CACGGAGGAG CTGATATTTG GTGGACA and AGCTTGTCCA CCAAATATCA GCTCCTCCGT GGATCTGCCT TGTCGAGCCT TCAGCTG were synthesized and purified by standard procedures. Annealing of these nucleotides into double-stranded structures left 4-base single stranded protruding ends compatible with those
10 generated by BamH I on one end of the molecule (GATC), and with Hind III-generated single stranded ends on the other end of the molecule (AGCT). Such annealed molecules were ligated into plasmid pBluescript SK(-) [hereinafter called pBSK; Stratagene Cloning Systems, La Jolla, CA], that had
15 been digested with BamH I and Hind III. The sequence of these oligonucleotides was such that when ligated onto the respective BamH I and Hind III sticky ends, the sequences of the respective recognition sites were maintained. An *E. coli* transformant harboring a plasmid containing the
20 oligonucleotide sequence was identified by restriction enzyme analysis, and the plasmid was named pMSV A.

Complementary oligonucleotides having the sequences AGCTGTGGAT AGGAGCAACC CTATCCCTAA TATACCAGCA CCACCAAGTC
AGGGCAATCC CGGG and TCGACCCGGG ATTGCCCTGA CTTGGTGGTG
25 CTGGTATATT AGGGATAGGG TTGCTCCTAT CCAC were synthesized and purified by standard procedures. Annealing of these nucleotides into double-stranded structures left 4-base sticky ends that were compatible with those generated by Hind III on one end of the molecule (AGCT), and with Sal
30 I-generated sticky ends on the other end of the molecule (TCGA). The sequence of these oligonucleotides was such that when ligated onto the Hind III sticky ends the recognition sequence for Hind III was destroyed.

DNA of pMSV A was digested with Hind III and Sal I, and was ligated to the above annealed oligonucleotides. An *E. coli* transformant harboring a plasmid containing the new oligonucleotides was identified by restriction enzyme site mapping, and was named pMSV AB.

Complementary oligonucleotides having the sequences CCGGGCCATT TGTTCCAGGC ACGGGATAAG CATTCAGCCA TGGGATATCA AGCTTGGATC CC and TCGAGGGATC CAAGCTTGAT ATCCCATGGC TGAATGCTTA TCCCGTGCCT GGAACAAATG GC were synthesized and purified by standard procedures. These oligonucleotides incorporate bases that comprise recognition sites for Nco I, EcoR V, Hind III, and BamH I. Annealing of these nucleotides into double-stranded structures left 4-base sticky ends that were compatible with those generated by Xma I on one end of the molecule (CCGG), and with Xho I-generated sticky ends on the other end of the molecule (TCGA). Such annealed molecules were ligated into pMSV AB DNA that had been digested with Xma I and Xho I. An *E. coli* transformant harboring a plasmid containing the oligonucleotide sequence was identified by restriction enzyme analysis, and DNA structure was verified by sequence analysis. The plasmid was named pMSV CPL. Together, these comprised the 5' untranslated leader sequence ("L") of the MSV coat protein ("CP") gene. These corresponded to nucleotides 167 to 186, and 188 to 317 of the MSV sequence of Mullineaux et al., (1984), and were flanked on the 5' end by the BamH I linker sequence GGATCCAG, and on the 3' end by the linker sequence GATATCAAGC TTGGATCCC. An A residue corresponding to base 187 of the wild type MSV sequence was inadvertently deleted during cloning.

The plasmid pMSV CPL DNA was digested at the Sma I site corresponding to base 277 of the MSV genomic

sequence, and the DNA was ligated to Bgl II linkers having the sequence CAGATCTG. An *E. coli* transformant harboring a plasmid having a unique Bgl II site at the position of the former Sma I site was identified and verified by DNA
5 sequence analysis, and the plasmid was named pCPL-Bgl.

The starting material in construction of a deleted version of the maize alcohol dehydrogenase 1 (Adh1) intron 1 is plasmid pVW119, which was obtained from V. Walbot, Stanford University, Stanford, CA. This plasmid contained
10 the DNA sequence of the maize Adh1.S gene, including intron 1, from nucleotides 119 to 672 [numbering of Dennis et al. (1984 Nucleic Acids Res. 12:3983-4000)], and was described in Callis et al. (1987 Genes and Develop.
1:1183-1200). In pVW119, the sequence following base 672
15 of Dennis et al. (1984) was GACGGATCC. The entire intron 1 sequence, with 14 bases of exon 1, and 9 bases of exon 2, was obtained from this plasmid on a 556 bp fragment following digestion with Bcl I and BamH I.

The plasmid pSG3525a(Pst) DNA was digested with BamH
20 I and Bcl I, and the 3430 bp fragment was purified from an agarose gel. DNA of plasmid pVW119 was digested with BamH I and Bcl I, and the gel purified fragment of 546 bp was ligated to the 3430 bp fragment. An *E. coli* transformant was identified that harbored a plasmid that generated
25 fragments of 3430 and 546 upon digestion with BamH I and Bcl I. This plasmid was named pSG AdhA1.

The DNA of pSG AdhA1 was digested with Hind III and with Stu I. The ends were made flush by T4 DNA polymerase treatment and then ligated. An *E. coli* transformant that
30 harbored a plasmid lacking Hind III and Stu I sites was identified and the DNA structure was verified by sequence analysis. The plasmid was named pSG AdhA1Δ. In this construct, 344 bp of DNA had been deleted from the

interior of the intron 1. The loss of these bases did not affect splicing of this intron. The functional intron sequences were obtained on a 213 bp fragment following digestion with Bcl I and BamH I.

5 DNA of plasmid pCPL-Bgl was digested with Bgl II, and the linearized DNA was ligated to the 213 bp Bcl I/BamH I fragment containing the deleted version of the Adh1.S intron 1 sequences from pSG AdhA1Æ. An *E. coli* transformant was identified by restriction enzyme site
10 mapping that harbored a plasmid that contained the intron sequences ligated into the Bgl II site, in the orientation such that the Bgl II/Bcl I juncture was nearest the 5' end of the MSV CPL leader sequence, and the Bgl II/ BamH I juncture was nearest the 3' end of the CPL. This
15 orientation was confirmed by DNA sequence analysis. The plasmid was named pCPL A1I1Æ. The MSV leader/intron sequences was obtained from this plasmid by digestion with BamH I and Nco I, and purification of the 373 bp fragment.

Construction of plant expression vectors based on the
20 enhanced 35S promoter, the MSV CPL, and the deleted version of the Adh1 intron 1 was as follows. DNA of plasmid p35S En²/NOS was digested with BamH I, and the 3562 bp linear fragment was ligated to a 171 bp fragment prepared from pMSV CPL DNA digested with BamH I. This
25 fragment contained the entire MSV CPL sequence. An *E. coli* transformant was identified by restriction enzyme site mapping that harbored a plasmid that contained these sequences in an orientation such that the Nco I site was positioned near the NOS Poly A sequences. This plasmid
30 was named p35S En² CPL/NOS. It contained the enhanced version of the 35S promoter directly contiguous to the MSV leader sequences such that the derived transcript included the MSV sequences in its 5' untranslated portion.

The DNA of plasmid pKA882 was digested with Hind III and Nco I, and the large 4778 bp fragment was ligated to an 802 bp Hind III/Nco I fragment containing the enhanced 35S promoter sequences and MSV leader sequences from p35S En² CPL/NOS. An *E. coli* transformant harboring a plasmid that contained fragments of 4778 and 802 bp following digestion with Hind III and Nco I was identified, and named pDAB310. In this plasmid, the enhanced version of the 35S promoter was used to control expression of the GUS gene. The 5' untranslated leader portion of the transcript contains the leader sequence of the MSV coat protein gene.

DNA of plasmid pDAB310 was digested with Nco I and Sac I. The large 3717 bp fragment was purified from an agarose gel and ligated to complementary synthetic oligonucleotides having the sequences CCGTACCTCGAGTTAAC and CATGGTTAACTCGAGGTACCGAGCT. These oligonucleotides, when annealed into double stranded structures, generated molecules having sticky ends compatible with those left by Sac I, on one end of the molecule, and with Nco I on the other end of the molecule. In addition to restoring the sequences of the recognition sites for these two enzymes, new sites were formed for the enzymes Kpn I (GGTACC), Xho I (CTCGAG), and Hpa I (GTTAAC). An *E. coli* transformant was identified that harbored a plasmid that contained sites for these enzymes, and the DNA structure was verified by sequence analysis. This plasmid was named pDAB1148.

DNA of plasmid pDAB1148 was digested with Bam HI and Nco I, the large 3577 bp fragment was purified from an agarose gel and ligated to a 373 bp fragment purified from pCPL A111E following digestion with Bam HI and Nco I. An *E. coli* transformant was identified that harbored a

plasmid that generated fragments of 3577 and 373 bp following digestion with BamH I and Nco I, and the plasmid was named pDAB303. This plasmid has the following DNA structure: beginning with the base after the final G residue of the Pst I site of pUC19 (base 435), and reading on the strand contiguous to the coding strand of the *lacZ* gene, the linker sequence ATCTGCATGG GTG, nucleotides 7093 to 7344 of CaMV DNA, the linker sequence CATCGATG, nucleotides 7093 to 7439 of CaMV, the linker sequence GGGGACTCTA GAGGATCCAG, nucleotides 167 to 186 of MSV, nucleotides 188 to 277 of MSV, a C residue followed by nucleotides 119 to 209 of *Adh1.S*, nucleotides 555 to 672 of maize *Adh1.S*, the linker sequence GACGGATCTG, nucleotides 278 to 317 of MSV, the polylinker sequence GTTAACTCGA GGTACCGAGC TCGAATTTC CC containing recognition sites for Hpa I, Xho I, Kpn I, and Sac I, nucleotides 1298 to 1554 of NOS, and a G residue followed by the rest of the pUC19 sequence (including the EcoR I site).

DNA of plasmid pDAB303 was digested with Nco I and Sac I, and the 3939 bp fragment was ligated to the 1866 bp fragment containing the GUS coding region prepared from similarly digested DNA of pKA882. The appropriate plasmid was identified by restriction enzyme site mapping, and was named pDAB305. This plasmid had the enhanced promoter, MSV leader and *Adh1* intron arrangement of pDAB303, positioned to control expression of the GUS gene.

The plasmid pDAB305 contained an enhanced 35S promoter with additional 3' sequences and embodied as nucleotides 7093 to 7344 of CaMV DNA, the linker sequence CATCGATG, nucleotides 7093 to 7439 of CaMV, the linker sequence GGGGACTCTA GAGGATCCAG, nucleotides 167 to 186 of MSV, nucleotides 188 to 277 of MSV, a C residue followed by nucleotides 120 to 210 of maize *Adh1.S*, nucleotides 555

to 672 of maize Adh1.S, the linker sequence GACGGATCTG, nucleotides 278 to 317 of MSV, and a G residue that represents the final base of an Nco I recognition sequence, CCATGG. As above, the GUS translational start
5 codon was part of the Nco I site. Transcripts from this promoter contain as the 5' untranslated leader essentially the MSV coat protein leader sequence, into which has been inserted a deleted version of the maize Adh1.S intron 1.

10

Claims

1. A method for transforming a plant cell which comprises:
 - (a) selecting at least one cell from a Type I, Type
5 II, hypocotyl-derived, or cotyledon-derived callus culture or from a meristem, pollen, cotyledon, or germ cell tissue; and
 - (b) inserting DNA into said cell by whisker mediated transformation.
- 10 2. A method for producing a fertile transgenic plant which comprises regenerating a cell produced by the method of claim 1.
3. A method for producing a fertile transgenic plant comprising the steps of: (i) establishing a regenerable
15 callus culture from a plant to be transformed wherein said callus culture is selected from the group consisting of Type I, Type II, hypocotyl-derived, and cotyledon-derived callus culture; (ii) selecting a plant cell aggregate therefrom for transformation; (iii) transforming said
20 plant cell aggregate with DNA by whisker-mediated transformation; (iv) identifying transformed cell lines therefrom; and (iv) regenerating fertile transgenic plants therefrom.
4. The method of Claim 3 wherein said Type I callus
25 culture is established from *Zea mays* or *Oryza sativa*.
5. The method of Claim 3 wherein said Type II callus culture is established from *Zea mays*.
6. The method of Claim 3 wherein said hypocotyl-derived culture is established from *Gossypium hirsutum*.
- 30 7. The method of Claim 3 wherein said cotyledon-derived culture is established from *Gossypium hirsutum*.

8. The method of Claim 3 wherein said plant cell aggregate is initiated on solid medium.
9. A fertile transgenic plant produced by the method of any one of claims 3-8.
- 5 10. A DNA construct functional in a plant cell comprising in the 5' to 3' direction of transcription, a transcriptional regulatory region functional in said plant cell and having a DNA sequence according to SEQ ID NO:1, and a gene of interest, said gene being either in the
10 sense or antisense orientation.
11. The transcriptional regulatory region according to Claim 10 comprising the DNA sequence SEQ ID NO:1.

SEQUENCE LISTING

<110> Petolino, Joseph F
Pareddy, Dayakar R
5 Hopkins, Nicole L
Armstrong, Katherine

<120> Whisker Mediated Transformation of Plant Cell
Aggregates and Plant Tissues and Regeneration of Plants
10

<130> 50518

<140>
<141>
15

<160> 1

<170> PatentIn Ver. 2.0

20 <210> 1
<211> 706
<212> DNA
<213> Agrobacterium tumefaciens

25 <400> 1
ctacaggcca aattcgctct tagccgtaca atattactca ccggtgcat gcccccac 60
gtaggtgaag gtggaaatta atgatccatc ttgagaccac aggcccacaa cagctaccag 120
tttcctcaag ggtccaccaa aaacgtaagc gcttacgtac atggtcgata agaaaaggca 180
atgtgtagat gttaacatcc aacgctcgctt tcagggatcc cgaattccaa gcttgggctg 240
30 cagggtcaatc ccattgcttt tgaagcagct caacattgat ctctttctcg agggagattt 300
ttcaaatacag tgcgcaagac gtgacgtaag tatccgagtc agtttttatt tttctactaa 360
tttggtcggt tatttcggcg tgtaggacat ggcaaccggg cctgaatttc gcgggtattc 420
tgtttctatt ccaacttttt cttgatccgc agccattaac gacttttgaa tagatacgt 480
gacacgcaa gcctcgctag tcaaaagtgt accaaacaac gctttacagc aagaacggaa 540
35 tgcgcgtgac gctcgcggtg acgccatttc gccttttcag aaatggataa atagccttgc 600
ttcctattat atcttcccaa attaccaata cattacacta gcatctgaat ttcataacca 660
atctcgatac accaaatcga ctctagaact agtggatccg tcgacc 706

40

INTERNATIONAL SEARCH REPORT

In tional Application No

PCT/US 99/01815

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/31 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 302 523 A (COFFEE RONALD ET AL) 12 April 1994 cited in the application see the whole document ---	1
X	WO 94 28148 A (ZENECA LTD) 8 December 1994 cited in the application see the whole document ---	1,2
X	NAGATANI N. ET AL.: "DNA delivery into rice cells and transformation using silicon carbide whiskers" BIOTECHNOLOGY TECHNIQUES, vol. 11, no. 7, 1997, pages 471-473, XP002103973 see the whole document --- -/--	1,2

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

27 May 1999

Date of mailing of the international search report

10/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

In tional Application No
PCT/US 99/01815

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 09696 A (PLANT GENETIC SYSTEMS NV) 11 June 1992 see the whole document ---	1-5, 8, 9
A	WO 97 12512 A (CALGENE INC) 10 April 1997 ---	1-3, 6-9
A	WANG K. ET AL.: "Whisker-mediated plant transformation: an alternative technology" IN VITRO CELLULAR AND DEVELOPMENTAL BIOLOGY, vol. 31, 1995, pages 101-104, XP002103974 see the whole document ---	1-9
A	SONGSTAD D. ET AL.: "Advances in alternative DNA delivery techniques" PLANT CELL, TISSUE AND ORGAN CULTURE, vol. 40, 1995, pages 1-15, XP002103975 see p.5 2.par - p.6 3.par. ---	1-9
A	WO 95 14098 A (CUI DECAI ; BIOTECHNOLOGY RES & DEV (US); NI MIN (US); GELVIN STANT) 26 May 1995 cited in the application see the whole document ---	10, 11
P, X	CHEMICAL ABSTRACTS, vol. 129, no. 8, 24 August 1998 Columbus, Ohio, US; abstract no. 91391, YAMAGUCHI, MASANORI ET AL: "Plant transformation method using crystal whiskers an improvement with DNA-basic substance complex" XP002103977 see abstract & JP 10 179174 A (HOKKO CHEMICAL INDUSTRY CO., LTD., JAPAN) ---	1-4, 9
P, X	SOUTHGATE E. ET AL.: "A comparison of methods for direct gene transfer into maize (Zea mays L.)" IN VITRO CELLULAR AND DEVELOPMENTAL BIOLOGY, vol. 34, July 1998, pages 218-224, XP002103976 see esp. p.217 1.col. 5.+6.par.; p.221 r.col. 2.+3.par.; table 4 -----	1-3, 5, 8, 9

INTERNATIONAL SEARCH REPORT

Information on patent family members

In tional Application No
PCT/US 99/01815

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5302523 A	12-04-1994	US 5464765 A	07-11-1995
WO 9428148 A	08-12-1994	AU 4339593 A	20-12-1994
WO 9209696 A	11-06-1992	AU 8914291 A	25-06-1992
		CA 2096843 A	24-05-1992
		EP 0558676 A	08-09-1993
		JP 6502533 T	24-03-1994
		US 5641664 A	24-06-1997
		US 5712135 A	27-01-1998
WO 9712512 A	10-04-1997	US 5846797 A	08-12-1998
		AU 7257996 A	28-04-1997
		CN 1198655 A	11-11-1998
		EP 0910239 A	28-04-1999
WO 9514098 A	26-05-1995	AU 687961 B	05-03-1998
		AU 1288595 A	06-06-1995
		BR 9408093 A	12-08-1997
		CN 1136825 A	27-11-1996
		EP 0729514 A	04-09-1996
		JP 9505205 T	27-05-1997

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.